

Characterization of Hilbert Spaces by the Strong Law of Large Numbers

JUN KAWABE

*Department of Information Sciences, Tokyo Institute of Technology,
Oh-okayama, Meguro-ku, Tokyo, Japan*

Communicated by the Editors

DEDICATED TO PROFESSOR HISAHARU UMEGAKI ON HIS 60TH BIRTHDAY

Hilbert spaces are characterized through the validity of the strong law of large numbers. Other characterizations of Hilbert spaces are also given at the same time in this note. © 1986 Academic Press, Inc.

1. INTRODUCTION

Let X be a real separable Banach space and X^* its topological dual space. Let $(\xi_n)_{n \geq 1}$ be a sequence of X -valued random variables defined on a basic probability space (Ω, \mathcal{A}, P) satisfying

$$E\xi_n = 0 \quad \text{for all } n \geq 1, \quad (1.1)$$

where E denotes the expectation (that is, Bochner integral with respect to P). Then (ξ_n) is said to *satisfy the strong law of large numbers* if

$$\frac{1}{n} \sum_{k=1}^n \xi_k \rightarrow 0 \quad \text{a. s.} \quad (= \text{almost surely}). \quad (1.2)$$

We shall then study under what conditions the strong law of large numbers holds. In [1] Beck has considered the condition

$$\sup_{n \geq 1} E\|\xi_n\|^2 < \infty \quad (1.3)$$

Received September 3, 1985; revised December 30, 1985.

AMS 1980 subject classification: Primary 60B12.

Key words and phrases: law of large numbers, covariance operators, nuclear operators, relative compactness, Banach space type.

and in [5] Hoffmann-Jørgensen and Pisier have considered the condition

$$\sum_{n=1}^{\infty} n^{-2} E \|\xi_n\|^2 < \infty. \quad (1.4)$$

In what follows, we denote by $L^2(X)$ the Banach space of all X -valued random variables which satisfy $E \|\xi\|^2 < \infty$ and, for each ξ in $L^2(X)$ denote by S_ξ its covariance operator (see [7]); that is, the nuclear operator of X^* into X which is defined by

$$S_\xi f = E \langle \xi, f \rangle \xi - \langle E \xi, f \rangle E \xi, \quad f \in X^*.$$

Further we denote by $\mathcal{N}(X^*, X)$ the Banach space of all nuclear operators of X^* into X with the nuclear norm $v(\cdot)$ (see [4]).

In this note we shall consider a condition

$$\sum_{n=1}^{\infty} n^{-2} S_{\xi_n} \quad \text{converges in } \mathcal{N}(X^*, X), \quad (1.5)$$

which is weaker than the above two conditions (1.3) and (1.4), and show that the strong law of large numbers holds for any sequence $(\xi_n)_{n \geq 1}$ of independent X -valued random variables satisfying (1.1) and (1.5) if and only if X is isomorphic to a Hilbert space. Other characterizations of Hilbert spaces are also given at the same time in the following theorem. Throughout this note, the letter R will denote the set of real numbers.

THEOREM. *For a real separable Banach space X , the following assertions (1)–(4) are equivalent:*

(1) *X is isomorphic to a Hilbert space.*

(2) *For any sequence $(\xi_n)_{n \geq 1} \subset L^2(X)$ satisfying (1.1), ξ_n converges in $L^2(X)$ if (and only if) the following two conditions are satisfied:*

(a) *For each $f \in X^*$, $\langle \xi_n, f \rangle$ converges in $L^2(R)$.*

(b) *The set $\{S_{\xi_n}\}$ is relatively compact in $\mathcal{N}(X^*, X)$.*

(3) *For any sequence $(\xi_n)_{n \geq 1}$ of independent X -valued random variables satisfying (1.1), $\sum_{n=1}^{\infty} \xi_n$ converges a. s. if*

$$\sum_{n=1}^{\infty} S_{\xi_n} \quad \text{converges in } \mathcal{N}(X^*, X). \quad (1.6)$$

(4) *The strong law of large numbers holds for any sequence $(\xi_n)_{n \geq 1}$ of independent X -valued random variables satisfying (1.1) and (1.5).*

2. AUXILIARY RESULTS

Before starting to prove the theorem, in this section we shall characterize relative compactness of sets and L^2 -convergence of sequences in $L^2(X)$ when X is a Hilbert space.

Denote by $l^p(X)$, $1 \leq p < \infty$, the Banach space of all sequences $x = (x_n)_{n \geq 1}$ of vectors in a Banach space X which satisfy $\sum_{n=1}^{\infty} \|x_n\|^p < \infty$. We shall use the following lemma to prove results in this section.

LEMMA 1 [2]. *A set $K \subset l^p(X)$ is relatively compact if and only if:*

(i) *For each $n \geq 1$, the set $K(n) = \{x_n; x \in K\}$ is relatively compact in X .*

(ii) $\lim_{N \rightarrow \infty} \sup_{x \in K} \sum_{n=N}^{\infty} \|x_n\|^p = 0$.

Let H be a real separable Hilbert space with the inner product (\cdot, \cdot) . Then $\mathcal{N}(X^*, X)$ coincides with the Banach space $\mathcal{T}(H)$ of all trace class operators on H .

PROPOSITION 2. *A set $K \subset L^2(H)$ is relatively compact if and only if the following conditions are satisfied:*

(1) *For each $x \in H$, the set $\{(\xi, x); \xi \in K\}$ is relatively compact in $L^2(R)$.*

(2) *The set $\{E\xi; \xi \in K\}$ is relatively compact in H and the set $\{S_\xi; \xi \in K\}$ is relatively compact in $\mathcal{T}(H)$.*

Proof. Assume first that K is relatively compact. Then condition (1) is satisfied since the map $\xi \rightarrow (\xi, x)$ of $L^2(H)$ into $L^2(R)$ is continuous. Condition (2) follows from an easily verified inequality,

$$v(S_\xi - S_\eta) \leq (E\|\xi - \eta\|^2)^{1/2} \cdot \{(E\|\xi\|^2)^{1/2} + (E\|\eta\|^2)^{1/2}\}$$

for all ξ, η in $L^2(H)$ satisfying $E\xi = E\eta = 0$.

Conversely, assume conditions (1) and (2). Let $\{\varphi_n\}$ be a basis of H and let us consider the set $M \equiv \{((\xi, \varphi_n))_{n \geq 1}; \xi \in K\}$ of elements in $l^2(L^2(R))$. Then M is relatively compact in $l^2(L^2(R))$, since it satisfies conditions (i) and (ii) of Lemma 1. In fact, condition (i) of Lemma 1 is satisfied from hypothesis (1). Since the map $x \rightarrow ((x, \varphi_n))_{n \geq 1}$ of H into $l^2(R)$ and the map $S \rightarrow ((S\varphi_n, \varphi_n))_{n \geq 1}$ of $\mathcal{T}(H)$ into $l^1(R)$ are continuous, it follows from hypothesis (2) that the set $\{((E\xi, \varphi_n))_{n \geq 1}; \xi \in K\}$ and the set $\{((S_\xi \varphi_n, \varphi_n))_{n \geq 1}; \xi \in K\}$ are relatively compact in $l^2(R)$ and $l^1(R)$, respectively.

Thus we have

$$\begin{aligned} & \lim_{N \rightarrow \infty} \sup_{\xi \in K} \sum_{n=N}^{\infty} E|(\xi, \varphi_n)|^2 \\ & \leq 2 \cdot \left\{ \lim_{N \rightarrow \infty} \sup_{\xi \in K} \sum_{n=N}^{\infty} |(E\xi, \varphi_n)|^2 + \lim_{N \rightarrow \infty} \sup_{\xi \in K} \sum_{n=N}^{\infty} (S\varphi_n, \varphi_n) \right\} = 0; \end{aligned}$$

therefore condition (ii) of Lemma 1 is also satisfied. We define now a continuous map of $l^2(L^2(R))$ into $L^2(H)$ as follows:

$$(\theta x)(\cdot) = \sum_{n=1}^{\infty} x_n(\cdot) \varphi_n, \quad x = (x_n(\cdot))_{n \geq 1} \in l^2(L^2(R)),$$

where the series is norm convergent in $L^2(H)$. Then we have $\theta M = K$ and consequently K itself is relatively compact in $L^2(H)$. From Proposition 2, using the standard argument, we have

COROLLARY 3. *A sequence $(\xi_n)_{n \geq 1} \subset L^2(H)$ converges in $L^2(H)$ if and only if the following conditions are satisfied:*

- (1) *For each $x \in H$, (ξ_n, x) converges in $L^2(R)$.*
- (2) *The set $\{E\xi_n\}$ is relatively compact in H and the set $\{S_{\xi_n}\}$ is relatively compact in $\mathcal{T}(H)$.*

3. PROOF OF THEOREM

(1) \Rightarrow (2) follows from Corollary 3. (2) \Rightarrow (3): Set $\zeta_n = \sum_{k=1}^n \xi_k$. It is sufficient to show that the sequence $(\zeta_n)_{n \geq 1}$ satisfies conditions (a) and (b) of (2) because $(\zeta_n)_{n \geq 1}$ converges a.s. if it converges in $L^2(X)$ (see [5, Lemma 1.2]). Condition (b) follows from (1.6) since $\sum_{k=1}^n S_{\xi_k} = S_{\zeta_n}$ for all $n \geq 1$. On the other hand, for each $f \in X^*$ we have

$$E \left| \sum_{k=m}^n \langle \xi_k, f \rangle \right|^2 = \sum_{k=m}^n \langle S_{\xi_k} f, f \rangle \leq v \left(\sum_{k=m}^n S_{\xi_k} \right) \cdot \|f\|^2$$

for all $n > m \geq 1$, and this, together with (1.6), implies condition (a). (3) \Rightarrow (4): By (1.1), (1.5), and assumption (3) we have that $\sum_{n=1}^{\infty} \xi_n/n$ converges a.s. since $\sum_{k=1}^n k^{-2} S_{\xi_k} = \sum_{k=1}^n S_{\xi_k/k}$ for all $n \geq 1$. Now we notice that Kronecker's lemma is valid in any Banach space (with the same proof as in the real case), so we have that $\sum_{k=1}^n \xi_k/n$ converges to 0 as $n \rightarrow \infty$. In order to prove (4) \Rightarrow (1) we need definitions of "type" and "cotype" and a result of [6].

Let $(\gamma_n)_{n \geq 1}$ be a sequence of independent standard Gaussian random variables. A Banach space X is said to be of *type 2* if for every sequence $(x_n)_{n \geq 1}$ in X such that $\sum_{n=1}^{\infty} \|x_n\|^2 < \infty$, we have that the series $\sum_{n=1}^{\infty} \gamma_n x_n$ converges a.s. and is said to be of *cotype 2* if for every sequence $(x_n)_{n \geq 1}$ such that the series $\sum_{n=1}^{\infty} \gamma_n x_n$ converges a.s., we have $\sum_{n=1}^{\infty} \|x_n\|^2 < \infty$. From a standard argument using the closed graph theorem we find that a Banach space X is of type 2 and of cotype 2 if and only if there exists a constant $C \geq 1$ such that for any positive integer n and any x_1, x_2, \dots, x_n in X ,

$$C^{-1} \sum_{k=1}^n \|x_k\|^2 \leq E \left\| \sum_{k=1}^n \gamma_k x_k \right\|^2 \leq C \sum_{k=1}^n \|x_k\|^2.$$

Then Kwapién has shown in [6, Proposition 3.1] that if X is of type 2 and of cotype 2 then it is isomorphic to a Hilbert space.

Now suppose that (4) holds, and we prove that if X is of type 2 and of cotype 2. First, since condition (1.5) is weaker than (1.4) it follows from Theorem 2.1 in [5] that X is of type 2. Next we prove that X is of cotype 2. Suppose to the contrary that there exists a sequence $(x_i)_{i \geq 1}$ such that $\sum_{i=1}^{\infty} \gamma_i x_i$ converges a.s., but $\sum_{i=1}^{\infty} \|x_i\|^2 = \infty$. (Without loss of generality we may assume that $x_i \neq 0$ for all $i \geq 1$.) If we set $a_k = \sum_{i=1}^k \|x_i\|^2$ then $a_k \rightarrow \infty$ and also $\sum_{k=1}^{\infty} \|x_k\|^2 / a_k = \infty$. Define a sequence $(\xi_k)_{k \geq 1}$ of independent X -valued random variables satisfying (1.1) with distributions such that

$$\Pr \left(\xi_k = -\frac{ka_k^{1/2} x_k}{\|x_k\|} \right) = \Pr \left(\xi_k = \frac{ka_k^{1/2} x_k}{\|x_k\|} \right) = \frac{1}{2} \cdot \frac{\|x_k\|^2}{a_k} \quad (3.1)$$

$$\Pr(\xi_k = 0) = 1 - \|x_k\|^2 / a_k.$$

Then by the Borel–Cantelli lemma, ξ_n/n does not converge to 0 a.s. so that

$$\frac{1}{n} \sum_{k=1}^n \xi_k \quad \text{does not converge to 0 a.s.} \quad (3.2)$$

Now let us consider a sequence $(\eta_n)_{n \geq 1}$ of X -valued Gaussian random variables defined by $\eta_n = \sum_{k=1}^n \gamma_k x_k$. Then, by (3.1) it is easy to show that $S_{\eta_n} = \sum_{k=1}^n k^{-2} S_{\xi_k}$ for all $n \geq 1$. Since η_n converges a.s., by Theorem 1 [3] $S_{\eta_n} = \sum_{k=1}^n k^{-2} S_{\xi_k}$ converges in $\mathcal{N}(X^*, X)$. Therefore from assumption (4) it follows that $\sum_{k=1}^n \xi_k/n$ converges to 0 a.s. and this contradicts (3.2). The proof is now complete.

ACKNOWLEDGMENTS

The author wishes to express his hearty thanks to Professor Hisaharu Umegaki for many kind suggestions and advice. The author also thanks the referee for his helpful comments.

REFERENCES

- [1] BECK, A. (1963). On the strong law of large numbers. In *Ergodic Theory* (F. B. Wright, Ed.), pp. 21–53. Academic Press, New York.
- [2] BROOKS, J. K., AND DINCULEANU, N. (1979). Conditional expectation and strong compactness in spaces of Bochner integrable functions. *J. Multivariate Anal.* **9** 420–427.
- [3] CHEVET, S. (1983). Gaussian measures and large deviations. In *Lecture Notes in Math.* Vol. 990, pp. 30–46. Springer-Verlag, Berlin/New York.
- [4] GROTHENDIECK, A. (1955). Produits tensoriels topologiques et espaces nucléaires. *Mem. Amer. Math. Soc.* **16** 1–191.
- [5] HOFFMAN-JØRGENSEN, J., AND PISIER, G. (1976). The law of large numbers and the central limit theorem in Banach spaces. *Ann. Probab.* **4** 587–599.
- [6] KWAPIÉN, S. (1972). Isomorphic characterizations of inner product spaces by orthogonal series with vector valued coefficients. *Studia Math.* **44** 583–595.
- [7] LINDE, V., TARIELADZE, V. I., AND CHOBANYAN, S. A. (1980). Characterization of certain classes of Banach spaces by properties of Gaussian measures. *Theory Probab. Appl.* **25** 159–164.